

## PATELLOFEMORAL PAIN SYNDROME (RUNNER'S KNEE)

### What is patellofemoral pain syndrome?

Patellofemoral pain syndrome is pain behind the kneecap. It has been given many names, including patellofemoral disorder, patellar malalignment, runner's knee, and chondromalacia.

### How does it occur?

Patellofemoral pain syndrome can occur from over-use of the knee in sports and activities such as running, walking, jumping, or bicycling.

The kneecap (patella) is attached to the large group of muscles in the thigh called the quadriceps. It is also attached to the shin bone by the patellar tendon. The kneecap fits into grooves in the end of the thigh bone (femur) called the femoral condyle. With repeated bending and straightening of the knee, you can irritate the inside surface of the kneecap and cause pain.

Patellofemoral pain syndrome also may result from the way your hips, legs, knees, or feet are aligned. This alignment problem can be caused by your having wide hips or underdeveloped thigh muscles, being knock-kneed, or having feet with arches that collapse when walking or running (a condition called over-pronation).

### What are the symptoms?

The main symptom is pain behind the kneecap. You may have pain when you walk, run, or sit for a long time. The pain is generally worse when walking downhill or down stairs. Your knee may swell at times. You may feel or hear snapping, popping, or grinding in the knee.

### How is it diagnosed?

Your healthcare provider will review your symptoms, examine your knee, and may order knee X-rays.

### How is it treated?

Treatment includes the following:

- Place an ice pack on your knee for 20 to 30 minutes every 3 to 4 hours for the first 2 to 3 days or until the pain goes away
- Elevate your knee by placing a pillow underneath your leg when your knee hurts.
- Take anti-inflammatory pain medicine, such as ibuprofen, as prescribed by your healthcare provider (adults aged 65 years and older should not take non-steroidal anti-inflammatory medicine for more than 7 days without their healthcare provider's approval)

- Do the exercises recommended by your healthcare provider or physical therapist.

Your healthcare provider may recommend that you:

- Wear shoe inserts (called orthotics) for over-pronation. You can buy orthotics at a pharmacy or athletic shoe store or they can be custom-made.
- Use an infrapatellar strap, a strap placed beneath the kneecap over the patellar tendon.
- Wear a neoprene knee sleeve, which will give support to your knee and patella.

While you are recovering from your injury, you will need to change your sport or activity to one that does not make your condition worse. For example, you may need to bicycle or swim instead of run. In cases of severe patellofemoral pain syndrome, surgery may be recommended. Your healthcare provider will show

### PATELLOFEMORAL PAIN SYNDROME (RUNNER'S KNEE)

